

Figure 1
Scheme For The Synthesis Of N-Methyl Piperazine




Figure 2A

Scheme A For The Synthesis Of N-Methyl Piperazine Acetic Acids




Figure 2B

Scheme B For The Synthesis Of N-Methyl Piperazine Acetic Acids




Figure 2C

Scheme C For The Synthesis Of N-Methyl Piperazine Acetic Acids




Figure 3A

Scheme A For The Synthesis Of ^{18}O Labeled N-Methyl Piperazine Acetic Acids




Figure 3B

Scheme B For The Synthesis Of ^{18}O Labeled N-Methyl Piperazine Acetic Acids




Figure 4A

Scheme A For The Synthesis Of Various Active Esters Of N-Methyl Piperazine
Via Imidazolid Formation




Figure 4B

Scheme B For The Synthesis Of Various Active Esters Of N-Methyl Piperazine
Via Oxallyl Chloride




Figure 4C

Scheme C For The Synthesis Of Various Active Esters Of N-Methyl Piperazine
Via Trifluoroacetate Ester




Figure 4D

Scheme For The Synthesis Of Various Active Esters Of N-Methyl Piperazine
Via Trifluoroacetate Esters




Figure 5A

Isotopic Pathway For Prepared N-Methyl Piperazine Acetic Acids





Figure 5B

Fragmentation of the Isobaric Label Set



NHS = N-hydroxysuccinimide

{} = Fragmentation Point


Stars indicate "heavy" isotopes

Pep = peptide

N* = ^{15}N ; C* = ^{13}C ; O* = ^{18}O


Figure 5C

Possible Fragment Ion Structures




Signature Ion 114

Or




Signature Ion 114




Signature Ion 115

Or




Signature Ion 115




Signature Ion 116

Or




Signature Ion 116



Signature Ion 117

Or



Signature Ion 117

Figure 6

General Fragmentation Properties Of The Set Of Isobaric Labeling Reagents

